




How much more could your lips say?

**Step 1:** Apply colored lipstick and kiss the card (do not pucker)



**Step 2:** Compare Lips

<b>Lips that appear disproportionate</b>	
	More fullness in the upper lip may create more pleasing proportions
<b>A lip border that could use greater definition</b>	
	Greater definition along the border of the upper lip may enhance the overall appearance of the lips
<b>Lips that are not as full or "pouty" as desired</b>	
	Adding fullness to both the upper and lower lip may produce the desired look

Restylane and Restylane-L can be used for lip enhancement in patients over 21 years and to add volume and fullness to the skin to correct moderate to severe facial wrinkles and folds, such as the lines from your nose to the corners of your mouth (nasolabial folds).

**Bring this card to your next appointment**

**Important Safety Considerations for Restylane and Restylane-L**

Restylane and Restylane-L should not be used by people with previous bad allergies, particularly to microorganisms known as gram-positive bacteria, or by people with serious allergies to drugs that have previously required in-hospital treatment. Restylane and Restylane-L should not be used by people with bleeding disorders. Restylane-L should not be used by anyone with a known allergy to lidocaine.

Use of Restylane or Restylane-L at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. In these instances, Restylane or Restylane-L use could delay healing or make your skin problems worse. After treatment, you might have some swelling, redness, pain, bruising, or tenderness. These are typically mild in severity and normally last less than 7 days in nasolabial folds and less than 14 days in lips. Swelling may be more likely in patients under 36 years, and bruising may be more likely in patients over 35 years. Rarely, the doctor may inject into a blood vessel, which can damage the skin. Although rare, red or swollen small bumps may occur. If you have had facial cold sores before, an injection can cause an outbreak. To avoid bruising and bleeding, you should not use Restylane or Restylane-L if you have recently used drugs that thin your blood or prevent clotting. Restylane and Restylane-L should not be used during pregnancy, when breastfeeding, or in patients under 18 years for nasolabial folds.

Treatment volume should be limited to 6.0 mL in wrinkles and folds, such as nasolabial folds, and limited to 1.5 mL per lip, as greater amounts significantly increase moderate and severe injection site reactions. The safety or effectiveness of treatment in areas other than nasolabial folds and lips has not been established in controlled clinical studies.

Restylane and Restylane-L are available only through a licensed practitioner. Complete product and safety information is available at [www.RestylaneUSA.com](http://www.RestylaneUSA.com).

Except as where otherwise indicated, all product names, slogans, and other marks are trademarks of the Valeant family of companies.